

## What Is a 4R Plant Nutrient?

By Tom Bruulsema, Plant Nutrition Canada; and Leanna Leverich Nigon, The Fertilizer Institute

November 15, 2023

Sustainability driven policies and business




Figure 1. The five interconnected aims of responsible plant nutrition and six key actions to take (SPRPN, 2020).

What is a plant nutrient? For many decades, plant nutrients have been defined by the concept of essentiality. More recently, however, 4R nutrient

stewardship has been defined as management that produces sustainable outcomes. Some elements provide benefits even if they are not essential. A new definition of what a plant nutrient is has been proposed and is under consideration by authorities in fertilizer regulation. Earn 1 CEU in Nutrient Management by reading this article and taking the quiz at https://web.sciencesocieties.org/Learning-Center/ Courses.

What is a plant nutrient? This question is important to 4R plant nutrition because it is important to include all plant foods that can be applied to improve the outcomes of nutrient stewardship.


For many decades, plant nutrients have been defined by the concept of essentiality. A mineral element was declared essential once science proved that in its absence, a plant could not complete its life cycle, and no other nutrient could substitute for it. In some jurisdictions, authorities required this proof of essentiality for nutrient claims in products sold as fertilizers.

More recently, however, 4R nutrient stewardship has been defined as management that produces sustainable outcomes, which suggests that it is the outcomes of applying the material, not its essentiality, that govern the questions, "What is the right source to apply?" and "What is a plant nutrient?" Some elements provide benefits even if they are not essential. Current classifications of essential and beneficial plant nutrient elements are outlined in Table 1.

Considering the foregoing, a new definition of what a plant nutrient is has been proposed and is under consideration by authorities in fertilizer regulation, including the Association of American Plant Food Control Officials (AAPFCO) and the International Standards Organization (ISO). The new definition proposes that "A mineral plant nutrient is an element which is essential or beneficial for plant growth and development or the quality attributes of the harvested product of a given plant species grown in its natural or cultivated environment" (Brown et al., 2022).

Whether a beneficial or essential nutrient, 4R nutrient stewardship practices should considered with every nutrient. The principles of 4R nutrient stewardship support the aims of responsible plant nutrition. Five such aims have recently been outlined, and they correspond with areas in which crop production can be made more sustainable (Figure 1; SPRPN, 2020). The five aims include: (1) improving productivity, income, efficiency, and resilience; (2) increasing nutrient recovery and recycling from waste; (3) lifting and sustaining soil health and soil carbon; (4) enhancing human health through nutrition[Bensitive agriculture; and (5) minimizing losses related to greenhouse gases, pollution of water and air, and biodiversity. Many nutrients contribute to achieving each aim, but some aims may need more particular nutrients. We will discuss each aim in turn and their dependence on nutrients.

### 1. Improving Productivity, Income, Efficiency, and Resilience



The first aim addresses productivity and its resulting benefits to income for growers while increasing efficiency and resilience. At least 30 to 50% of crop yield is attributable to commercial fertilizer nutrient inputs (Stewart et al., 2005). This means that the efficiency of crop production per unit of land and per unit

Figure 2. The "Law of the Minimum" states that if any essential nutrient is deficient, then crop growth will be reduced even if all of the other nutrients are adequately supplied (Carl Sprengel and Justus von Liebig; International Plant Nutrition Institute, 2019).

of other fixed inputs is increased by 40 to 100%. Net income can thus be more than doubled, provided fertilizers are applied using the right source, right rate, right timing, and the right placement. The nutrients most frequently limiting yield are nitrogen (N), phosphorus (P), and potassium (K) with sulfur (S) and micronutrients increasingly playing important roles as well (Flis & Jones, 2020).

The "Law of the Minimum" means that any nutrient can limit yield and the use efficiency of other nutrients (Figure 2). Sound diagnosis of the soil and past crops is an important part of any 4R nutrient plan. This means continued testing to determine whether any nutrient may contribute to yield improvement. The new definition of plant nutrient opens greater opportunity for adaptive research and adaptive management within 4R programs.

It's not easy to find hard numbers for resilience, but it seems logical to argue that creating balanced nutrient levels in soils contributes to improving it. University and Extension typically encourage growers to maintain soils at a "medium" or optimum soil test level for P and K and other nutrients. At an optimum level, the soil provides most of the nutrient to the crop, and the purpose of fertilizing is replenishing the soil rather than directly boosting yield. There is thus more flexibility in "right rate," and growers can adjust for changing prices of fertilizers and crops as well as suffer less yield loss if

weather conditions prevent timely application.

Building balanced nutrient levels will lead to healthy, resilient soils and greater productivity and profitability for growers. Both essential and beneficial nutrients can contribute to this balance under the 4R nutrient stewardship framework.

#### 2. Increasing Nutrient Recovery and Recycling From Waste

As the human population expands, the depletion of the resources used to make fertilizers and the accumulation of wastes in soils and the natural environment have become important concerns. Both concerns can be mitigated by recycling where possible. The biggest opportunity for N, P, and K is by improving the management of manure. As indicated by the nutrient balances in NuGIS (TFI, 2023), the differences between the amounts excreted by livestock and the recovered amounts applied to cropland are large. Since manures are bulky and difficult to transport, technologies to extract nutrients in forms similar to fertilizer are needed to improve the overall utilization of the nutrients.

There are also opportunities to recapture N further down the value chain, by capturing it in wastewater treatment to produce biosolids. Most of these wastes that contain N and P also contain other important nutrients, including S, and micronutrients, especially zinc (Zn). "High quality biosolids and similar materials may be used beneficially in sustainable agriculture" (Chaney, 1994) if risks of excesses of arsenic, cadmium, copper, molybdenum, and selenium are managed (Chaney, 2012).

Recovery of struvite from wastewater is increasingly being used to produce an ammonium magnesium phosphate fertilizer with an analysis of 5½8½000Mg in a granular form suitable for blending with other common fertilizers. In Pennsylvania,

centrifuge extraction of P from dairy manure was shown to be economically feasible if the other nutrients extracted and environmental benefits were valued (Rotz et al., 2022).

The various nutrient extraction technologies facilitate increased recycling of nutrients—both essential and beneficial—within the agriffood system, and we can expect to see more of them in the future.



Improving the management of manure presents a huge opportunity for increasing nutrient recovery of N, P, and K. Photo courtesy of Camille Resseguier, INRAE.

## 3. Lifting and Sustaining Soil Health and Soil Carbon

The use of 4R practices to manage nutrients is critical to support crop photosynthesis and make soil carbon storage an effective proposition for greenhouse gas mitigation. What is required is a delicate balancing act. Increasing primary productivity on cropland, reducing wastes, selecting climates mart nutrient sources, and using

inhibitors to manage nitrous oxide emissions are all critical. Nitrogen inputs in particular play a large role in increasing soil organic carbon in the temperate soils of North America (Alvarez, 2005). Phosphorus can also be important. A recent metall analysis of 116 peerlieviewed papers showed that the addition of P fertilizer generally increases total N in soils with higher biological retention, lower N losses, and increased soil carbon levels (Wang, 2022). The metalanalysis shows the important link among P, N, and C in maintaining resilient and healthy soils.

The strong role of N and other nutrients in the multiple mechanisms of soil carbon storage underscores the need for integrated consideration of 4R nutrient management in programs that address both the emissions and sinks associated with cropping systems while keeping them productive (Bruulsema, 2022). Improving the carbon footprint of crop production requires simultaneous attention to three areas at once: soil carbon, nitrous oxide emission, and N use efficiency.

Beyond soil carbon, nutrient inputs play an important role in replenishing soil fertility, another important aspect of soil health. The status of soil pH, P, K, Mg, S, Zn, and Cl in North America is monitored through soil testing and has been reported on systematically every five years or so since 2001 (TFI, 2021b). Soil has capacity to retain nutrients, and regular soil testing helps ensure that soil fertility is maintained at an optimum level, important for both soil health and the resiliency of crop production systems.

## 4. Enhancing Human Health Through Nutrition Sensitive Agriculture

Nutrients enhance not only yield, but nutritional quality as well. Most plant nutrients are nutrients for animals and humans as well. The role of several micronutrients is particularly important for the alleviation of hidden hunger, which affects more than 2

billion people globally. Hidden hunger is most prevalent in the African continent and South Asia but is also increasingly reported in Europe and in North America. Children and women of reproductive age are most vulnerable (Stevens et al., 2022). Seven mineral nutrients (iron, zinc, copper, calcium, magnesium, selenium, and iodine) and several vitamins are often lacking in human diets (Van Der Straeten et al., 2020). Selenium and iodine have been added to fertilizers used in Finland and China, respectively, to improve human nutrition, even though these elements are not considered essential to plants and only rarely benefit crop yields.

Plant[based foods provide 56% of the protein intake and 40–85% of various minerals to the global population (Smith et al., 2022). The remainder is supplied largely from products from animals that, in turn, were largely dependent upon plants for nutrition. So, getting the mineral nutrients into harvested crops is important. A global metall analysis of 67 studies found that fertilizing corn with Zn for yield improvement raised Zn concentration from 22 to 27 ppm, a 25% increase, but not enough to meet the 38 ppm target for alleviating human Zn deficiency (Mutambu at al., 2023).

Atmospheric CO<sub>2</sub> is predicted to surpass 550 ppm in the next 30–80 years. Many food crops grown at 550 ppm show reductions of 3–17% in protein and Zn contents compared with current conditions. This could cause an additional 175 million people to be Zn deficient and an additional 122 million people to be protein deficient by the middle of this century (Smith & Myers, 2018). Crop nutrition strategies may need to change as a result.

Crop advisers need to pay attention to crop quality attributes in the assessment of 4R practices implemented onlifarm. Selection of appropriate cultivars and inclusion of micronutrients in applied fertilizers can enhance the concentration of these nutrients in harvested crops.

# 5. Minimizing Losses Related to Greenhouse Gases, Pollution of Water and Air, and Biodiversity




Figure 3. Trends in N outputs and inputs for harvested crops in North America from 1961 to 2018. Outputs do not include forage crops. Inputs include fertilizer, manure, legume fixation, and atmospheric deposition. The slope of a line from any point to the origin indicates N use efficiency. Figure reprinted from Bruulsema (2021; Data sources: Zhang et al., 2021, with extrapolation from 2015 to 2018 using FAO, 2020).

When it comes to impacts on the environment, two nutrients dominate the discussion. Phosphorus losses have been long known to impact water quality. Nitrogen losses affect not only water quality, but also air quality and greenhouse gas emissions. Improvement of nutrient use efficiency (NUE) goes a long way toward reducing these losses (SPRPN, 2023). Trends in NUE for North American crop production are improving or at least keeping up with increases in crop yields (as indicated in NuGIS in TFI [2021a] and in the cropland nutrient budgets of FAO [2022]), but further stewardship efforts are needed to keep nutrient surpluses to a minimum in intensified crop production.

The aggregate performance for North American crop production has encouraging trends. For the past few decades, both N use efficiency and N output per acre have been increasing (Figure 3). But some important caveats need to be kept in mind (Bruulsema, 2021):

 The large role of soybeans in the N balance hides the lower efficiency of other crops. As a Nfixing species, soybeans and other legumes have high N use efficiency.

- Corn grain is lower in protein today compared with decades ago.
- As yields increase, higher levels of N use efficiency are needed to limit N surplus.

Biological N fixation involves cobalt, nickel, molybdenum, and sometimes vanadium, and thus these are required in greater amounts in NIfixing species. Nickel is required in greater amounts when urea or ammonia are the dominant N source; Si is highly beneficial when Mn is present at toxic levels (Brown et al., 2022).

The challenge to continue improvements in N use efficiency is still relevant. It requires attention to all aspects of crop production, including 4R plant nutrition that includes consideration of all plant nutrients, essential and beneficial, for their potential to contribute.

#### Take Home Message

So, what is a 4R plant nutrient? It's any nutrient that enhances plant growth and development or the quality attributes of the harvested crop products. These nutrients are all included in the aims of responsible plant nutrition. Paying attention to the nutrient composition of the harvested crop may be an important part of a 4R nutrient management plan, particularly when purchasers of crop commodities offer incentives.

#### References

AAPFCO. (2019). *Product label guide*. Association of American Plant Food Control Officials. https://www.rs.uky.edu/regulatory/fertilizer/label\_guide\_2019.pdf Alvarez, R. (2005). A review of nitrogen fertilizer and conservation tillage effects on soil organic carbon storage. *Soil Use and Management*, 21(1), 38–52. https://doi.org/10.1111/j.14750

Brown, P.H., Zhao, F.D., & Dobermann, A. (2022). What is a plant nutrient? Changing definitions to advance science and innovation in plant nutrition. Plant and Soil, 476(1), 11–23. https://doi.org/10.1007/s11104ID21ID5171Iw

Bruulsema, T. (2021). Nutrient stewardship: Taking 4R further. *Crops & Soils Magazine*, 55(1), 34–40. https://doi.org/10.1002/crso.20165

Bruulsema, T. (2022). Soil carbon needs 4R nutrients. Crops & Soils Magazine, 55(4), 34–38. https://doi.org/10.1002/crso.20203

Chaney R.L. (1994). Trace metal movement: Soil@lant systems and bioavailability of biosolids@applied metals. In C.E. Clapp, W.E. Larson, and R.H. Dowdy (Eds.), Sewage sludge: land utilization and the environment (pp. 27–31). SSSA. Chaney, R.L. (2012). Chapter two—Food safety issues for mineral and organic fertilizers. Advances in Agronomy, 117, 51–116. https://doi.org/10.1016/B978@012@94278@4.00002@

EU. (2019). Regulation (EU) 2019/1009 of the European Parliament and of the Council of 5 June 2019 laying down rules on the making available on the market of EU fertilising products. https://eurllex.europa.eu/homepage.html

FAO. (2022). FAOSTAT and IFA cropland nutrient budget database. http://www.fao.org/faostat/en/#data/ESB

Flis, S., & Jones, J. (2020). Sulfur 4R management and the changing landscape of fertility. Crops & Soils Magazine, 53(6), 19–22. https://doi.org/10.1002/crso.20074 Havlin, J., Tisdale, S., Nelson, W., & Beaton, J. (2013). Soil fertility and fertilizers: An introduction to nutrient management (8th edition). International Plant Nutrition

Institute. (2019). Soil fertility manual. The Fertilizer Institute. ISO. (2022). International standards. Fertilizers, soil conditioners and beneficial substances—classification (7851) and vocabulary (8157). International Organization for Standardization. https://www.iso.org/obp/ui#home

Mutambu, D., Kihara, J., Mucheru[Muna, M., Bolo, P., & Kinyua, M. (2023). Maize grain yield and grain zinc concentration response to zinc fertilization: A meta@analysis. Heliyon, 9(5). https://doi.org/10.1016/j.heliyon.2023.e16040

Rotz, C.A., Reiner, M.R., Fishel, S.K., & Church, C.D. (2022). Whole farm performance of centrifuge extraction of phosphorus from dairy manure. Applied Engineering in Agriculture, 38(2), 321–330. https://doi.org/10.13031/aea.14863

SPRPN. (2020). A new paradigm for plant nutrition (Issue Brief 01). Scientific Panel on Responsible Plant Nutrition. https://www.sprpn.org/post/alhewlparadigmlforlplantll nutrition

SPRPN. (2023). Defining nutrient use efficiency in responsible plant nutrition (Issue Brief O4). Scientific Panel on Responsible Plant Nutrition. https://sprpn.org/issuellbrief/defininglhutrientluselefficiencyllnlresponsiblelplantlhutrition

Smith, M.R., & Myers, S.S. (2018). Impact of anthropogenic CO<sub>2</sub> emissions on global human nutrition. Nature Climate Change, 8(9), 834–839. https://doi.org/10.1038/s41558

Smith, N.W., Fletcher, A.J., Millard, P., Hill, J.P., & McNabb, W.C. (2022). Estimating cropland requirements for global food system scenario modeling. Frontiers in Sustainable Food Systems, 6, https://doi.org/10.3389/fsufs.2022.1063419

Stevens G.A., Beal, T., Mbuya, M.N.N., Luo, H., & Neufeld, L.M. (2022). Micronutrient deficiencies among preschool@ged children and women of reproductive age worldwide: a pooled analysis of individual@evel data from population@representative surveys. *The Lancet Global Health*, e1590–e1599. https://doi.org/10.1016/S22140 109X(22)00367D

Stewart, W.M., Dibb, D.W., Johnston, A.E., & Smyth, T.J. (2005). The contribution of commercial fertilizer nutrients to food production. Agronomy Journal, 97(1), 1–6. https://doi.org/10.2134/agronj2005.0001

TFI. (2021a). NuGIS Nutrient Use Geographic Information System. The Fertilizer Institute. https://nugis.tfi.org TFI. (2021b). Soil test levels in North America, 2020 summary update. The Fertilizer Institute. https://soiltest.tfi.org

Van Der Straeten, D., Bhullar, N.K., De Steur, H., Gruissem, W., MacKenzie, D., Pfeiffer, W., ... & Bouis, H. (2020). Multiplying the efficiency and impact of biofortification through metabolic engineering. Nature Communications, 11(1), 5203. https://doi.org/10.1038/s41467\( \tilde{D}\) 20\( \tilde{D}\) 20\( \tilde{D}\)

Wang, R., Bicharanloo, B., Hou, E., Jiang, Y., & Dijkstra, F.A. (2022). Phosphorus supply increases nitrogen transformation rates and retention in soil: A global metalanalysis. *Earth's Future*, *10*, e2021EF002479. https://doi.org/10.1029/2021EF002479

## **Self-Study CEU Quiz**

Earn 1 CEU in Sustainability by taking the quiz for the article at https://web.sciencesocieties.org/Learning-Center/Courses. For your

convenience, the quiz is printed below. The CEU can be purchased individually, or you can access as part of your Online Classroom Subscription.

1. <b>V</b>	/hich of	the fo	ollowing	is trı	ue of	the	static	chamber	method?
-------------	----------	--------	----------	--------	-------	-----	--------	---------	---------

- a. Gas samples are collected manually.
- b. Gas concentrations are measured in the lab.
- c. Gas concentrations are measured with a gas chromatograph.
- d. All of the above.
- 2. The Dairy Soil and Water Regeneration project aims to capture the environmental footprint of feed practices on poultry farms.
  - a. True.
  - b. False.
- The DSWR project began its work in 2021 with a survey of dairies in \_\_
   major milk production regions of the country.
  - a. two
  - b. three
  - c. four
  - d. six
- 4. What does "dynamic" and "static" refer to when describing the dynamic or static measurement system?
  - a. If a researcher has to manually measure the sample in the lab.
  - b. If the air is or is not circulating during the measurement.
  - c. If the chamber can or cannot be moved around the field.
  - d. Whether or not the chamber has a lid.

	Carbon Enhancement Network."
	a. True.
	b. False.
6.	What data are researchers in the DSWR project collecting with
	infiltrometers?
	a. Field-saturated hydraulic conductivity.
	b. Soil temperature.
	c. Air temperature.
	d. Nitrous oxide emissions.
7.	For the DSWR project, what is a typical daily change in oxygen levels in
	intermediate or well-drained soils?
	a. 9-11%.
	b. 10-20%.
	c. 19-21%
	d. 15-30%
8.	Why might researchers be missing some measurements for GHG
	emissions during the winter?
	a. They are using instruments that cannot operate in winter temperatures.
	b. They are missing GHG releases that occur during freeze-thaw events.
	c. They are only measuring CO🏻 and not CH🗂 or N🖾.
	d. They are using instruments that cannot operate in precipitation.
9.	Feedback from DWSR researchers is helping LI-COR update both the
	instrument's hardware and software.
	a. True.
	b. False.

5. GRACEnet stands for "Greenhouse Gas Reduction Through Agricultural

- 10. The data loggers used by DWSR to measure temperature, moisture, and oxygen do not allow the researchers to remotely access these data.a. True.
  - b. False.

Text © . The authors. CC BY-NC-ND 4.0. Except where otherwise noted, images are subject to copyright. Any reuse without express permission from the copyright owner is prohibited.