

## Nutrient Loss Trade-Offs Between Conventional and No-Till Systems

January 12, 2024



A cover-cropped field with snow and ice cover in Chatham, Ontario. Photo by Ryan Carlow.

Nutrient losses through assisted "tile" drains from agricultural fields contribute to algal blooms in surface waters globally. Noltill is a practice that can reduce particulate phosphorus (PP) losses from fields, but this can increase dissolved reactive phosphorus (DRP) losses in tile drainage. It is unclear if cover ("catch") crops can offset some of these issues.

Researchers investigated tile drain discharge from a conservation[fill field along Lake Erie in Ontario (considered conventional for the region) and an adjacent no[fill field incorporating cover crops. They compared seasonal and annual water yields and concentrations and losses of phosphorus and nitrogen between the two site types and tracked soil fertility and management activity for four years.

The researchers found that while water losses from the systems were similar, losses of sediments, PP, and nitrate were greater from the conservation till site. Losses of DRP were greater from the notitil, covertropped site, offsetting the lower PP losses such that there was little difference in total phosphorus losses between sites, largely due to incidental losses following surface application of fertilizer. This study provides insight into tradeloffs in nutrient losses from tile drainage using different management systems.

## Adapted from

Macrae, M. L., Plach, J. M., Carlow, R., Little, C., Jarvie, H. P., McKague, K., Pluer, W. T., & Joosse, P. (2023). Tradelbffs in nutrient and sediment losses in tile drainage from nollill versus conventional conservation conservation systems. *Journal of Environmental Quality*, *52*, 1011–1023. https://doi.org/10.1002/jeq2.20502

Text © . The authors. CC BY-NC-ND 4.0. Except where otherwise noted, images are subject to copyright. Any reuse without express permission from the copyright owner is prohibited.